

TIME-RELATED FACTORS IN RESEARCH ON DIET AND CANCER

DEMETRIUS ALBANES, ARTHUR SCHATZKIN and MARC S. MICOZZI

Cancer Prevention Studies Branch, Prevention Program, Division of Cancer Prevention and Control, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892, U.S.A.

Abstract—Time-related factors have been an integral aspect of laboratory and epidemiologic studies concerning the effects of diet on cancer. In this survey, we illustrate several ways in which consideration of time factors has furthered our understanding in this area. As one of the key dimensions in descriptive studies of secular trends and migrant populations, time factors suggest general associations between diet and cancer. The investigation of time-related parameters such as age, duration of dietary exposure, and time from exposure to cancer, lends greater specificity to the diet-cancer relationship. Both micro- and macro-nutrients are examined, as well as nutrition-mediated factors such as growth and anthropometry. Time-related issues relevant to the design of future observational and intervention studies of diet and cancer (i.e. critical etiologic periods, timing of dietary assessment, biological indicators, and secular trends) are also discussed.

Anthropometry Cancer Diet Epidemiology Nutrition Time

INTRODUCTION

Research on diet and cancer has become an important scientific focus in recent years [1], although the view of diet as an etiologic factor in human carcinogenesis is not new [2]. The role played in the development of cancer by dietary macronutrients such as fats, fiber, and total calories, as well as by several vitamins (e.g. A, C, and E) and minerals (e.g. selenium) is under intense investigation. Time-related factors have been an important aspect of both laboratory and epidemiologic studies in this area. As one of the key dimensions in descriptive studies, time factors (along with person and place) have suggested *general* relationships between dietary factors and cancer. The investigation of time-related factors such as age, duration of dietary exposure, and time from exposure to cancer outcome, lends greater specificity to the diet-cancer relationship. A broad and, necessarily, schematic overview of these issues is presented, drawing examples from the available literature. In addition, the relevance of such time considerations to the design of epi-

demologic studies and cancer chemoprevention trials is examined.

TEMPORAL ASPECTS IN DIET-CANCER RESEARCH

The diet-cancer relation has been explored in a number of descriptive studies that examine secular trends, international correlations and migrant populations. While these studies have contributed to our current understanding by providing evidence for general associations between diet and cancer, they usually do not lend specificity to the suggested relations.

Secular trends

Changes in cancer incidence or mortality over time have been related to fluctuations in dietary factors in an effort to explore possible diet-cancer associations. For example, Hems [3] assessed secular trends of dietary components and breast cancer mortality for the United Kingdom taking place between 1911 and 1971. Some of these findings are illustrated in Fig. 1. The trend for breast cancer mortality is seen to




Fig. 1. Change with time of breast cancer mortality among women of England and Wales, and fat, animal protein and sugar in the diet (g/day) from 1911-15 to 1971-75 [3].

parallel those for dietary fat, animal protein, and sugar intake over the six decades shown. These data, coupled with supporting experimental data and several plausible mechanisms (e.g. enhanced estrogen metabolism or bile acid production), suggest a role for dietary fat in the etiology of breast cancer. The increase in breast cancer mortality beginning around 1960 apparently occurred 10-20 years after an increasing trend in dietary intake: the correlation between regional breast cancer rates and dietary fat was higher when intake of 10 years earlier was used (correlation coefficient $r = 0.6$), than when either the concurrent diet ($r = 0.4$) or that of two decades earlier ($r = 0.1$) was used. In contrast, international correlation analyses we have conducted using food availability data for fat and other nutrients, and 1973-1977 incidence rates for various cancers, show little or no effect of lag-time [4]. Correlating incidence data [5] from 1 year with either concurrent food data or those from approximately 12 years earlier [6] for 21 countries, we have observed for several cancer sites little difference in the correlations between the food data from either period and cancer incidence. For example, the Pearson correlation coefficient for breast, prostate, and colon cancers are 0.71, 0.56, and 0.79, respectively, when concurrent dietary fat data are used. For the food data from 12 years prior to the year of cancer incidence, the coefficients are 0.75, 0.59, and 0.79, respectively. Thus, while such studies cannot make inferences about the individual, they are supportive of associations such as between breast cancer and dietary fat.

Migrant studies

Descriptive studies of migrants have also contributed to our hypotheses concerning the effect of diet on cancer. Migrant populations are observed to undergo changes in exposure to suspected environmental risk factors (including diet) and health outcomes through time and space, while genetic factors may be considered relatively constant. Acculturation, the process by which traditional lifestyle is substituted over time for host country patterns, provides a necessary link between migration and dietary change. Determining the actual time of acculturation is not easily done, however, as the year of migration, as well as the age at migration, may independently affect the year of acculturation.

Several investigations have related the incidence of various neoplasms among Japanese migrants to their degree of acculturation to a Western lifestyle and diet. One such etiologic study of stomach cancer among male Japanese migrants in Hawaii by Haenszel *et al.* [7] found that migrants (but not their offspring) exhibited increased risk comparable to that of men in Japan, and that this elevated risk was associated with the consumption of specific Japanese foods such as pickled vegetables and salted fish, among others. In another study of migrants, Buell [8] describes the increasing incidence of breast cancer in successive generations of Japanese women in America, as compared to Japan (Table 1). He suggests that these changing rates may be attributable to dietary acculturation involving increased macronutrient intake, and reflected, at least in part, in increased height and weight during the growth period. However, data on specific dietary changes were not presented, so that the precise association with diet remains unclear in this study.

In contrast, Nomura *et al.* [9] compared the diet of Japanese men whose wives had developed breast cancer with the diets of other Japanese men who had participated in the Japan-Hawaii Cancer Study (the authors assumed that the diets of the husbands and wives were similar). Their results indicated that husbands of the breast cancer cases consumed a Western-style diet (more beef or other meat,

Table 1. Age-adjusted breast cancer incidence rates

Age	Japan	First generation	Second generation	U.S. whites
35-64	31.8	93.6	116.5	179.4
65-74	21.8	163.4	—	293.4

butter/
Japanes
group.
changes
populat

Anthrop

Study
human
variable
changes
pometri
ing pas
macron
effects

Anth
related
quency
Americ
in the
Americ
and 19
with th
same an
have ra
whites.
Japanes
older Ja
the low
in cancer
the yo
women
and stat
the olde
These g
ger (and




Fig. 2. Age-adjusted breast cancer incidence rates among Japanese men and women.

migrants have also
 eses concerning the
 rant populations are
 ages in exposure to
 risk factors (including
 s through time and
 s may be considered
 uration, the process
 le is substituted over
 ns, provides a neces-
 and dietary change.
 e of acculturation is
 as the year of mi-
 e at migration, may
 ear of acculturation.
 ave related the inci-
 ns among Japanese
 of acculturation to a
 One such etiologic
 among male Japanese
 enszel *et al.* [7] found
 r offspring) exhibited
 e to that of men in
 d risk was associated
 ecific Japanese foods
 es and salted fish,
 study of migrants,
 reasing incidence of
 e generations of Ja-
 ca, as compared to
 sts that these chang-
 ble to dietary accul-
 ed macronutrient in-
 in part, in increased
 the growth period.
 dietary changes were
 e precise association
 in this study.

al. [9] compared the
 ose wives had devel-
 the diets of other
 participated in the
 y (the authors as-
 husbands and wives
 s indicated that hus-
 er cases consumed a
 beef or other meat,

cancer incidence rates	
Second generation	U.S. whites
116.5	179.4
—	293.4

butter/margarine/cheese, and corn), and ate less Japanese foods, than did men in the control group. Further examination of specific dietary changes occurring over time among migrant populations would be useful.

Anthropometry

Studying the nutritional component of human growth patterns and anthropometric variables offers an alternative assessment of changes in dietary factors over time. Anthropometry not only provides information regarding past nutritional history (particularly for macronutrients), but can evaluate age-specific effects of nutrition [10].

Anthropometric changes over time have been related to the increase in breast cancer frequency across successive cohorts of Japanese-American women [11]. This increase can be seen in the age-specific cancer rates for Japanese-American women in Los Angeles between 1972 and 1977. These rates are contrasted in Fig. 2 with the incidence for whites residing in the same area. Younger Japanese-American women have rates that are similar to those of U.S. whites, and are much higher than those of old Japanese-American women; the rates of the older Japanese-American women are similar to the low rates among women in Japan. Changes in cancer rates parallel growth patterns in that the younger cohort of Japanese-American women (born in 1955) evidenced greater growth and stature at age 17 years (158 cm; [12]) than the older (born in 1940) cohort (154 cm; [13]). These growth differences suggest that the younger (and taller) cohort had longer intake of a

more Western diet which may have contributed to their increased incidence of breast cancer.

Given that growth is a function of dietary intake, international correlations between height or other anthropometric measures at different ages and cancer incidence or mortality can provide additional information about the general impact of diet, particularly over-nutrition, on breast and other cancers. We studied the relationships between age-specific anthropometric measurements [14], and breast cancer incidence [5] and mortality [15] rates in 34 populations around the world. Table 2 shows the correlation between breast cancer mortality and age-specific stature, weight and biacromial diameter. The correlation between breast cancer and each of these anthropometric variables increases with age across childhood and adolescence, and is highest for the oldest age groups. Similar results were obtained for breast cancer incidence [16]. Since dietary differences across countries are better reflected in growth differences at later rather than earlier ages, the fact that these correlations with breast cancer are stronger for later ages offers further support for the connection between diet and cancer.

Timing of dietary exposure

Time-related factors can also lend greater specificity to the nature of exposure (in this case diet), and allow more precise delineation of the association between diet and cancer than do descriptive studies. These parameters include: (1) age at exposure, (2) the duration of exposure, and (3) the time from exposure to tumor or cancer outcome. Unfortunately, few data are currently available from human studies which shed light on these issues with regard to diet-cancer associations. Laboratory experi-




Fig. 2. Age-specific breast cancer incidence in Japanese vs white females, Los Angeles, U.S., 1972-1977 [5].

Table 2. Cross-national correlation of age-specific anthropometric measures with age-adjusted breast cancer mortality

Age	Stature	Weight	Biacromial width
6	0.48	0.48	0.30
7	0.38	0.39	0.29
8	0.56	0.40	0.44
9	0.56	0.42	0.40
10	0.53	0.37	0.44
11	0.45	0.37	0.27
12	0.57	0.42	0.31
13	0.65	0.43	0.32
14	0.62	0.50	0.41
15	0.75	0.58	0.47
16	0.76	0.69	0.68
17	0.74	0.68	0.78
18	0.72	0.75	0.71

ments, however, have contributed to our knowledge in this area.

For example, in a series of studies involving comprehensive monitoring of diet, body size, and tumorigenesis in rats, Ross *et al.* [17] observed quantitative, time-dependent relationships between nutrition, growth, body size, and neoplasia. Both the age at various dietary exposures and the duration of those intake patterns were found to be highly predictive of subsequent tumorigenesis. Specifically, these investigators demonstrated that increased tumor incidence in animals was associated with: (1) a rapid growth rate during early postnatal life, (2) high absolute protein intake after weaning, (3) high efficiency of converting consumed food into body mass at the time of puberty, (4) high protein intake relative to body weight in early adulthood, and (5) high levels of food intake during adulthood.

Numerous experiments such as those of Verma *et al.* [18] and Tannenbaum and Silverstone [19] involve micronutrient supplementation (e.g. vitamin A derivatives) or dietary manipulation (e.g. fat or total calorie restriction) at various ages or periods of carcinogenesis. Although generalizations to humans from animal experiments must be made cautiously, these and similar studies suggest that chemopreventative agents or dietary modification in humans may have anticarcinogenic effects, and that the timing of such interventions may be important. These investigations have shown, for example, that dietary factors can operate during the promotion stage of carcinogenesis, long after initiation has occurred [18, 20]. Thus, laboratory models afford unique opportunities to generate hypotheses which can be further tested through epidemiologic research.

RESEARCH DESIGN ISSUES

Time-related factors have important implications for the design of diet and cancer studies in humans. Four research aspects which require the attention of investigators include: (1) the evaluation of *critical etiologic periods*, (2) methods of *dietary assessment*, (3) utilization of *biologic indicators* of nutritional status, and (4) the effects of *secular trends*.

Critical etiologic periods

Although seldom addressed, each of the time-related factors of age, duration, and the time

from exposure to cancer endpoints, should be considered when planning human diet-cancer research. In addition, hypotheses concerning multistage carcinogenesis in humans and the effects of diet on that process may suggest which periods of exposure should be investigated. For example, if dietary fat is believed to exhibit both early and late stage effects on breast carcinogenesis in animals [19, 21] and in humans [22], then case-control and cohort studies of this cancer should attempt to collect data for both early and late exposures. In prospective studies specifically, the time of data collection and the hypotheses concerning duration of exposure and latency of its effect will determine how long the investigators wait for study endpoints.

With regard to chemoprevention trials, time factors relating to the intervention are critical to study design. The primary intervention in this case is nutrient supplementation or dietary modification in a controlled, randomized fashion. Generally, agents believed to be effective in later stages of carcinogenesis are being employed in adult study populations [23]. Nutritional manipulation having primarily early-stage effects would necessitate the use of younger study populations for much longer periods in order to adequately test the efficacy of the intervention. Decisions must be made regarding the appropriate age and duration of intake for study participants, the lag-time of effect, and the pre-clinical latency period of the cancer, in order to exclude, or analyze separately, cancers present at the start of the study. Also, age-dependent competing causes of illness or death should be considered, since a large number of non-cancer deaths occurring among subjects can impede subsequent interpretation of cancer endpoints.

Dietary assessment

Epidemiologic studies in this area are hampered by difficulties in accurately assessing dietary intake. The several methods of assessing current diet that have been developed differ, in part, with respect to the time period over which intake is measured. The 24-hour recall, for example, includes all foods eaten over the previous day, while the food frequency instrument is designed to elicit average consumption of various foods over a longer period (e.g. 1 year). Block [24] has reviewed advantages and disadvantages of the various methods of relevance to current epidemiologic research. A particular

time-re
metho
in mic
ation
miscla
made
Sempo
and he
the nu
to red

As e
intake
inform
past, e
Howev
the ini
clinical
past r
conne
during
[27, 28
the dis
partic
biased
Byers
tion o
gave
dietary
practic
past e
maine
import
cancer
ranted
tive st
availa
Invest
spent
or co
vestig
from
about
where

Biolo

Sim
data
odolo
sures
Blood
nutrit
plorel
prosp
strate

endpoints, should be human diet-cancer hypotheses concerning humans and the may suggest which be investigated. For eved to exhibit both on breast carcinoma and in humans [22], short studies of this collect data for both prospective studies a collection and the ration of exposure determine how long study endpoints. evention trials, time ention are critical to intervention in this entation or dietary d, randomized fash- ved to be effective in genesis are being y populations [23]. having primarily ecessitate the use of s for much longer tely test the efficacy ions must be made age and duration of ents, the lag-time of atency period of the clude, or analyze t at the start of the competing causes of considered, since a er deaths occurring de subsequent inter-nts.

in this area are accurately assessing methods of assessing n developed differ, in ne period over which 24-hour recall, for s eaten over the pre- frequency instrument age consumption of r period (e.g. 1 year). dvantages and disad- thods of relevance to search. A particular

time-related problem with the 24-hour recall method is the marked intra-individual variation in micronutrient consumption. Day-to-day variation can lead to severe dietary exposure misclassification when comparisons are being made among individuals. Liu *et al.* [25] and Sempos *et al.* [26] have examined this problem and have estimated for several micronutrients the number of 24-hour dietary recalls necessary to reduce misclassification to a specified level.

As difficult as it is to assess current dietary intake, it is more problematic to obtain reliable information on what a person ate in the distant past, especially in childhood and adolescence. However, because of the long period between the initiation of the carcinogenic process and its clinical manifestation, what people ate in the past may be as relevant to the diet-cancer connection as what they are eating now or during the previous few years. Earlier studies [27, 28] concluded that dietary information from the distant past tended to be unreliable, and, in particular, that estimates of past diet were biased by current intake. A recent study by Byers *et al.* [29], though, suggests that recollection of dietary practices from the distant past gave a better estimate of originally recorded dietary data than did the assessment of current practices, although the biased recollection of past dietary practices by the current diet remained a problem [24]. Given the potentially important influence of early diet on the risk of cancer, further research in this area is warranted. Re-interviews of subjects from prospective studies on whom earlier dietary data are available may be helpful in this regard [30]. Investigations of dietary recall in persons who spent early years in institutions such as schools or convents may also be instructive. Investigators may also obtain current dietary data from study participants and obtain information about past diet from the parents of participants, where available.

Biologic indicators

Since collection of valid early dietary intake data is constrained by a number of methodologic difficulties, the use of alternative measures of diet and nutrition becomes imperative. Blood nutrient levels offer such measures of nutritional status, which have been widely explored in diet-cancer studies. For example, a prospective study by Kark *et al.* [31] demonstrated that persons developing cancer had

significantly lower serum vitamin A (i.e. retinol) levels at least 12 months before diagnosis than did persons remaining free from cancer (41.3 vs 46.9 $\mu\text{g}/100\text{ ml}$). Assessing vitamin A stability at -20°C , and that during periods of freeze-thaw and exposure to light (simulating conditions which may have occurred during the 14-year follow-up period), the authors showed no degradation of samples tested. While these and other investigators excluded cancers which developed soon (e.g. 1 year) after serum collection in an effort to minimize the effects of disease on nutrition, others specifically evaluate serum levels from cases which develop at different times throughout the surveillance period [32]. Thus, while the effects of cancer on such parameters can complicate the interpretation of case-control studies which obtain blood from known cases, prospective collection of appropriately stored blood samples may offer more meaningful tests for hypotheses.

Other useful nutritional markers include tissue samples such as toenails, which can be analyzed for selenium or other materials, and adipose tissue (reflecting dietary fatty acid composition), which may offer assessment of more distant nutritional status advantageous in various research settings. Also, as discussed earlier, anthropometry offers valid estimates of childhood, adolescent, or adult intake of important macronutrients by measuring parameters such as height, weight or skinfold thickness in adults which are enduring indicators of dietary intake [10].

Secular trends

Trends in diet and other variables should also be considered in the design of prospective studies and clinical trials concerned with diet and cancer, since secular trends can affect the results of these investigations. For example, the U.S. food supply has undergone changes since the beginning of this century, including increases in animal protein, total fat, cholesterol, vitamins A and C, as well as some of the B vitamins [33]. In contrast, consumption of crude fiber and carbohydrates has decreased, especially that from cereal products and potatoes. In addition, the use of vitamin and/or mineral supplements among adults 55-64 years old has increased in the U.S. from 34% in 1971, to 40% in 1980 [34]. Observations such as these should be considered when planning study sample size, subject recruitment, and other aspects of study design.

REFERENCES

1. Committee on Diet, Nutrition and Cancer, Assembly of Life Sciences, National Research Council. *Diet, Nutrition and Cancer*. Washington, D.C.: National Academy Press; 1982.
2. Hoffman FL. *The Mortality from Cancer throughout the World*. Newark, New Jersey: The Prudential Press; 1915.
3. Hems G. Associations between breast-cancer mortality rates, child-bearing and diet in the United Kingdom. *Br J Cancer* 1980; 41: 429-436.
4. Albanes D. Unpublished data. 1984.
5. Waterhouse J, Muir C, Shanmugaratnam K, Powell J. *Cancer Incidence in Five Continents*, Vol. IV. Lyons, France: IARC Scientific Publication No. 42, 1982.
6. Food and Agriculture Organization: *Food Balance Sheets and Per Capita Food Supplies, 1961-1977*. Rome: United Nations; 1980.
7. Haenszel W, Kurihara M, Segi M, Lee RKC. Stomach cancer among Japanese in Hawaii. *J Natl Cancer Inst* 1972; 49: 969-988.
8. Buell P. Changing incidence of breast cancer in Japanese-American women. *J Natl Cancer Inst* 1973; 51: 1479-1483.
9. Nomura A, Henderson BE, Lee J. Breast cancer and diet among the Japanese in Hawaii. *Am J Clin Nutr* 1978; 31: 2020-2025.
10. Johnston FE. *Anthropometry and Nutritional Status*. Committee on Food Consumption Patterns. Food and Nutritional Board, National Research Council. Washington, D.C.: National Academy Press; 1981: 252-264.
11. Micozzi MS. Nutrition, body size and breast cancer. *Yearbook of Phys Anthropol* 1985; 28: 175-206.
12. Greulich WW. A comparison of the physical growth and development of American-born and native Japanese children. *Am J Phys Anthropol* 1957; 15: 489.
13. Kondo S, Eto M. Physical growth studies on Japanese-American children in comparison with native Japanese. In *Proc Meeting for Review and Seminar of the U.S.-Japan Cooperative Research on Human Adaptabilities*. Kyoto: Society for the Promotion of Science, and National Science Foundation, 1972.
14. Eveleth PB, Tanner JM. *Worldwide Variations in Human Growth*. Cambridge: Cambridge University Press; 1976.
15. Kurihara M, Aoki K, Tominaga S. *Cancer Mortality Statistics in the World*. Nagoya, Japan: University of Nagoya Press; 1984.
16. Micozzi MS, Schatzkin A. International correlation of anthropometric variables and adolescent growth patterns with breast cancer incidence. *Am J Phys Anthropol* 1985; 66: 206-207.
17. Ross MH, Lustbader ED, Bras G. Dietary practices of early life and spontaneous tumors of the rat. *Nutr Cancer* 1982; 3: 150-167.
18. Verma AK, Shapas BG, Rice HM, Boutwell RK. Correlation of the inhibition by retinoids of tumor promoter-induced mouse epidermal ornithine decarboxylase activity and of skin tumor promotion. *Cancer Res* 1979; 39: 419-425.
19. Tannenbaum A, Silverstone H. Nutrition in relation to cancer. In: Greenstein JB, Jaddow A, Eds. *Advances in Cancer Research*, Vol. 1. New York: Academic Press; 1953: 451-501.
20. Tannenbaum A. The dependence of the genesis of induced skin tumors on the caloric intake during different stages of carcinogenesis. *Cancer Res* 1944; 4: 673-677.
21. Carroll KK, Khor HT. Effects of dietary fat and dose level of DMBA on mammary tumor incidence in rats. *Cancer Res* 1970; 30: 2260-2264.
22. Miller AB. Role of nutrition in the etiology of breast cancer. *Cancer* 1977; 39: 2704-2708.
23. Micozzi MS, Boone CW, Kelloff GJ, Tangrea JA, Helzlsouer KJ, Taylor PR. Chemoprevention of cancer: implications for clinical pharmacology. *J Clin Pharm* 1985; 25: 164-175.
24. Block G. A review of validations of dietary assessment methods. *Am J Epidemiol* 1982; 115: 492-505.
25. Liu K, Stamler J, Doyer A *et al.* Statistical methods to assess and minimize the role of intra-individual variability in obscuring the relationship between dietary lipids and serum cholesterol. *J Chron Dis* 1978; 31: 399-418.
26. Sempos CT, Johnson NE, Smith EL, Gilligan C. Effects of intraindividual and interindividual variation in repeated dietary records. *Am J Epidemiol* 1985; 121: 120-130.
27. Acheson ED, Doll R. Dietary factors in carcinoma of the stomach: a study of 100 cases and 200 controls. *Gut* 1964; 5: 121-131.
28. Gregor A, Lee PN, Roe JC, Wilson MJ, Melton A. Comparison of dietary histories in lung cancer cases and controls with special reference to vitamin A. *Nutr Cancer* 1980; 2: 93-97.
29. Byers TE, Rosenthal RI, Marshall JR, Rzepka TF, Cummings KM, Graham S. Dietary history from the distant past: a methodological study. *Nutr Cancer* 1983; 5: 69-77.
30. Jensen OM, Wahrendorf J, Rosenqvist A, Geser A. The reliability of questionnaire-derived historical dietary information and temporal stability of food habits in individuals. *Am J Epidemiol* 1984; 120: 281-290.
31. Kark JD, Smith AH, Switzer BR, Hames CG. Serum vitamin A (retinol) and cancer incidence in Evans County, Georgia. *J Natl Cancer Inst* 1981; 66: 7-16.
32. Nomura AMY, Stemmermann GN, Heilbrun LK, Salkeld RM, Vuilleumier JP. Serum vitamin levels and the risk of cancer of specific sites in Hawaiian males of Japanese ancestry. *Cancer Res* 1985; 45: 2369-2372.
33. Gortner WA. Nutrition in the United States 1900 to 1974. *Cancer Res* 1975; 35: 3246-3253.
34. National Center for Health Statistics: Unpublished data. 1985.

The var
exposur
how we
what h
relevan
a partic
was ca
tempor
because
anisms
type of
tends t
another
regardi
presum
probabl
pend up
age.

The
caused
though
great d
both ex